

Quantum quairesonances in grazing incident angle atom-surface collisions

Antonia Ruiz,¹ José P. Palao,² and Eric J. Heller³

¹*Departamento de Física Fundamental y Experimental,
Universidad de La Laguna, La Laguna 38204, Spain*

²*Departamento de Física Fundamental II, Universidad de La Laguna, La Laguna 38204, Spain*

³*Departments of Chemistry and Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

(Dated: June 24, 2008)

The momentum transfer between the normal components to an index direction in the collision of an atom with a periodic surface is investigated. For fast atoms with grazing angle of incidence there is an interval of azimuthal angles around the index direction for which the energy transfer can be very efficient. This effect is reflected in quantum diffraction patterns with large non-specular peaks, associated with the parallel to the surface and normal to the index direction momentum component, and can be described in terms of quairesonance. Although the classical dynamics does not reproduce the precise quantum diffraction probabilities, indicating the quantum nature of this effect, classical and quantum computations show that the span of the quairesonance region coincides in both dynamics and can be classically estimated from the phase-space analysis.

PACS numbers:

Recent experiments in grazing angle of incidence collisions of fast atoms with a surface along a low index direction have shown quantum diffraction patterns even for a total translational energy as large as several keV [1, 2, 3]. This result is explained by the existence of two motion regimes, a fast one along a low index direction, with momentum $p_x = \sqrt{2mE} \sin \theta \cos \varphi$, and a slow one in the plane perpendicular to the index direction, with momentum components $p_y = \sqrt{2mE} \sin \theta \sin \varphi$ and $p_z = \sqrt{2mE} \cos \theta$, where E is the total incident energy and m the atom mass. The component p_z is chosen normal to the surface. For incidence along an index direction, $\varphi_i = 0$ and $p_{yi} = 0$. Within these conditions, the fast atoms “feel” the surface as structureless parallel furrows aligned along the index direction [1], which constitutes an example of channelling on surfaces [2, 4]. As a consequence the two regimes of motion are decoupled, resulting in a constant momentum component p_x . Under grazing angle conditions the energy associated with the normal components is very small, and, as the change in p_y is quantized, a quantum diffraction pattern is observed despite the relatively large total translational energy.

However, the previous analysis does not explain to what extent the diffraction pattern still survives for incidence not in a surface index direction ($p_{yi} \neq 0$), nor does it predict how efficient is the energy transfer between the normal components responsible for the diffraction pattern. A convenient analysis of the issue can be made in terms of quairesonance. The concept of quairesonance was first introduced to explain some very efficient and specific energy transfer between the vibrational and rotational degrees of freedom experimentally observed in state resolved collisions between vibrotationally excited diatom molecules and one atom [5, 6]. An striking observation was a remarkable insensitivity of the quairesonant effect to the specific details of the interaction potential

[5, 7, 8]. It was also shown that a fully classical treatment of the collision reproduces the experimental results remarkably well [6, 7, 9, 10, 11]. Recently, the quairesonance analysis has been extended to new processes in different classical systems [12, 13], including grazing angle atom surface collisions [14].

An important open question is whether the quantum effects in the dynamics share any common mechanism with the classical quairesonances. This issue has been analyzed in ultracold atom-diatom collisions [10]. The results were not conclusive as for the low energies considered the main inelastic quantum channels associated with quairesonances remain closed. Grazing angle atom-surface collisions in a close-to an index incident direction provides a new ground to study the similarities between both dynamics.

Classically, quairesonance can be understood in terms of the adiabatic invariance theory and the method of averaging [15]. Let us consider a two-dimensional integrable system described by the action variables (J_1, J_2) , perturbed by a interaction, depending on other additional degrees of freedom, which is switched on and off. The interaction induces a coupling between the two “internal” degrees of the perturbed system, and defines a process from an initial state (J_{1i}, J_{2i}) to a final state (J_{1f}, J_{2f}) . Let us assume (a) a slowly varying perturbative interaction; and (b) an initial internal state satisfying the approximate $M : N$ non-linear resonance condition $M\omega_{2i} - N\omega_{1i} \approx 0$, with M and N small integers, and ω_α the two independent frequencies of the system. The analysis of dynamics of the nearly resonant system in terms of the standard secular perturbation theory introduces a canonical transformation to new action variables, $I_1 = -J_1/N, I_2 = J_2 + MJ_1/N$, defined in a rotating frame in which the angle variable ϕ_2 associated with the action I_2 oscillates much more rapidly than the other variables during the entire transient process. The aver-

aging of the transformed Hamiltonian over this rapidly oscillating angle leads to its conjugated action as the first term of the series expansion for the adiabatic invariant of the system in the proximity of the nonlinear resonance. In terms of the original actions, the existence of this local adiabatic invariance implies that

$$\frac{\Delta J_2}{\Delta J_1} = -\frac{M}{N}, \quad (1)$$

where $\Delta J = J_f - J_i$. This accurate correlation between the action changes is the central result of the quasisresonance analysis [12, 13]. The quasisresonance region is defined as the interval of initial values for which the conditions (a) and (b) are satisfied.

In atom surface collisions, classical quasisresonance applies to an effective two-dimensional free atom, described by the components of the momentum parallel to the surface, which is perturbed by the transient atom-surface interaction. Here the analysis can be simplified considering a coordinate system with the x-axis fixed along the index direction of interest, since in this coordinate system the actions I_1 and I_2 can be directly identified with the components of the momentum p_y and p_x . The “external” translational action that controls the strength of the interaction is p_z .

Assumption (a) corresponds to a collision with large translational energy (perturbative regime) and a small p_z (slowly variant interaction); both together implying $p_z \ll p_x$. Meanwhile the assumption (b) corresponds to $p_y \ll p_x$. These conditions define a scattering process with a grazing incident beam closely aligned to an index direction of the surface. Since $\dot{x} \gg \dot{y}, \dot{z}$, the averaging over this rapidly oscillating variable ($\phi_2 \equiv x$) allows us to identify its conjugated action variable, the momentum along such direction ($I_2 \equiv p_x$), as the first order adiabatic invariant characterizing quasisresonance. The dynamics in the perpendicular plane (yz) is dictated by the Hamiltonian

$$H_{av} = \frac{1}{2m} (p_y^2 + p_z^2) + V_{av}(y, z) \quad (2)$$

with the averaged potential

$$V_{av}(y, z) = \frac{1}{L} \int_0^L V(x, y, z) dx, \quad (3)$$

being L the surface periodicity along the x-axis. In a quasisresonance region, the coupling induced by this averaged potential enables an energy transfer between the normal components of the momentum, p_y and p_z . The efficiency of such energy transfer is determined by the form of V_{av} resulting from the original potential and the particular index direction implied. In general, the energy transfer is more efficient for incidence close to a low index direction. Outside quasisresonance such energy transfer does not occurs, and incident beams “feel” a plain surface and only suffer an effective specular reflection.

In a coordinate system with the x-axis fixed along the surface direction $(1, 0)$, the adiabatic invariance of the momentum along a symmetry direction (M, N) , associated with the approximated resonance condition $Mp'_y - Np'_x \simeq 0$, leads to the correlation $\Delta p'_y / \Delta p'_x = -M/N$ [14]. In this coordinate system the efficient energy transfer in the plane perpendicular to the direction (M, N) is reflected by significant changes in both internal actions, p'_x and p'_y . This is the usual result of the general quasisresonance analysis.

Quasisresonance is satisfied for grazing incident beams aligned over an interval of azimuthal angles around the crystal index direction. We define the width W_{py} of the quasisresonance region as the interval of initial values of p_y for which the approximated resonance condition (b) is satisfied, or, alternatively, W_φ as the corresponding interval of incident azimuthal angles φ_i . Both quantities are related by $W_\varphi = 2 \sin^{-1}[W_{py}/(2\sqrt{2mE} \sin \theta_i)]$. W_{py} can be estimated from the analysis of the phase space associated with the averaged Hamiltonian at frozen values z_{fr} of the normal z coordinate [13]. We focus on the dynamics around the resonant incident direction, $p_{yi} = 0$, which can be properly described by the pendulum like Hamiltonian $H_{pend} = p_y^2/2m + V_{av}(y, z_{fr})$. The maximum excursion in action of the islands of the resonance zone in this integrable system provides the simple expression

$$W_{py} \approx 2\sqrt{2m[V_{av}(y_u, z_m) - V_{av}(y_s, z_m)]}, \quad (4)$$

being z_m the normal coordinate that gives the maximum strength interaction at incident energy E , y_u the hyperbolic point associated with the innermost separatrix, and y_s the most stable fixed point of the potential $V_{av}(y, z_m)$. The expression (4) is valid provided that the resonance zones associated with the different crystal index directions remain well isolated in phase space. In a non perturbative regime, the overlapping between neighbouring resonance zones in phase space makes the quasisresonance regions shrink [13].

To illustrate quasisresonance we simulate the atom-surface collision considering a Lennard-Jones and Devonshire potential with parameters taken from Ref. [16], describing the elastic scattering of He atoms from a perfectly periodic rigid LiF(001) surface. The low normal velocities implicit under grazing angle conditions, and the robustness of the quasisresonance phenomena with respect to the details of the interaction potential justify the use of this simple model.

Figure 1 shows the changes in p_x and p_y for classical trajectories closely aligned to an index direction. Two different behaviors are clearly distinguished; there are significant changes in p_y in the region around the symmetry direction $\varphi_i = 0$, identified as the quasisresonance region, and a constant value of p_y outside it. The adiabatic invariant in the quasisresonance p_x remains constant in the whole depicted range. The change in p_z (not shown




FIG. 1: (Color online) $\Delta k_x = (p_{xf} - p_{xi})/\hbar$ (black dashed-dotted line) and Δk_y (brown dots) versus the initial azimuthal angle φ_i , for classical trajectories closely aligned to the [110] direction ($\varphi_i = 0$). All the trajectories have incident polar angle $\theta_i = 0.506\pi$, initial $z_i = 60\text{\AA}$ and (x_i, y_i) chosen at random in a unit cell. The red solid line and the blue dashed line represent the mean values and the root-mean-square values of Δk_y respectively, obtained at each angle φ_i from an average over 2000 classical trajectories with identical initial conditions, except for the coordinates (x_i, y_i) . The mean values (red squares) and the root-mean-square values (blue circles) derived from the quantum diffraction probabilities are also depicted.

in the figure) is dictated by the energy conservation in the global elastic process.

For the quasisresonance conditions (a) and (b), the quantum mechanical treatment of the collision can be simplified, as only a small number of quantum channels are effectively open [1]. Under these conditions the large momentum component along the index direction remains constant and only quantum transitions corresponding to reciprocal lattice vectors perpendicular to this direction present a non-negligible diffraction probability. As only a small amount of energy associated with the perpendicular degrees of freedom is available, only a few quantum channels contribute to the diffraction spectrum. This effect has been experimentally confirmed for incidence along a low crystal index [1, 2, 3]. Then the quantum dynamics can be described by an effective 2D model [1] with the averaged potential Eq. (3).

Figure 2 shows the diffraction spectra for four incident azimuthal angles: along a crystal index direction, inside, on the edge, and outside the classical quasisresonance region. The significant non-specular diffraction peaks in the first three spectra correspond with the efficient energy transfer between the perpendicular momentum components, p_y and p_z , predicted in the classical quasisresonance. As the incident direction approaches the edge the intensity of the inelastic channels decreases, be-




FIG. 2: Quantum (solid lines) and quasiclassical (dashed lines) diffraction spectra for a grazing incident beam with energy $E = 200\text{ eV}$, polar angle $\theta_i = 0.506\pi$ and different azimuthal angles φ_i ; (a) along the [110] direction, $\varphi_i = 0$, (b) inside the quasisresonance region, $\varphi_i = 0.01$, (c) on the edge of the quasisresonance region, $\varphi_i = 0.02$, and (d) outside of the quasisresonance region, $\varphi_i = 0.04$. The quasiclassical probabilities were calculated replacing the continuous distribution by an histogram. The diffraction probabilities have been convoluted with a Lorentzian function to account for possible broadening effects.

ing the specular diffraction peak the only that survives for incidence outside the quasisresonance region. The precise distribution of the diffraction probabilities is dictated by quantum effects in the dynamics of the system, and as expected, is not correctly reproduced by the classical model, see figure 2. The important result however is the qualitative agreement, and especially the correct prediction of a significant qualitative change in the quantum diffraction spectrum as the incident direction crosses the border of quasisresonance. Both the exact classical and exact quantum results confirm the simple quasisresonance analysis. As figure 1 shows, there is also a very good agreement between the trends in classical and quantum changes in p_y . Hence both classical and quantum dynamics reflect the same mechanism of efficient energy transfer that characterizes quasisresonance around the low index direction.

Figure 3 shows the dependence of the span of the quasisresonance region on the total translational energy. The monotonic increase of the width W_{p_y} with the energy is a peculiar feature of the simple interaction potential considered. Since in this model the averaged potential is zero except for the principal surface directions (1, 0) and (0, 1), quasisresonance around other symmetry directions is a higher order perturbative effect. Therefore, the main mechanism for the eventual demise of the main quasisresonance regions, i.e., the overlapping between neighbouring resonance zones in phase space, is nearly absent. The




FIG. 3: The span of the quiresonance region around the [110] direction versus the incident energy E , in a process with grazing incident angle $\theta_i = 0.506\pi$. Labels on the left correspond the angular width W_φ ; the squares are the numerical classical estimation, the crosses the numerical quantum estimation, and the solid line the classical prediction derived from (4). Labels on the right give the size of the quiresonance region in momentum space, $W_{ky} = W_{py}/\hbar$; the dashed line corresponds to the classical estimation (4). As a criterion for the numerical estimation of the angular width of the quiresonance region, we consider the full width at half height of root-mean-square values of Δk_y .

angular with W_φ decreases rapidly at low energies and changes very slowly at higher energies. At low energies, no normal quantum inelastic channels are energetically allowed and, as expected, there is not good agreement between the quantum and classical dynamics. Both results become nearly identical as the energy increases and more diffraction channels are open.

In conclusion, the grazing angle collision of an atom with a periodic surface for incidence closely aligned to a (low) index direction has been investigated using both a classical and a quantum description. For a large interval of incident energies the results can be analyzed in terms of quiresonance. This interval is limited below by low energies for which the dynamics is not perturbative, and above by large energies with large normal to the surface component of the momentum for which the atom-surface interaction is not slowly variant. Quiresonance is characterized in this system by the adiabatic invariance of the momentum component along a symmetry direction of the target surface, together with an efficient energy transfer between the normal degrees of freedom. Even when the classical results do not reproduce the precise quantum diffraction probabilities, the averaged results are in good agreement in both dynamics, specially for large energies. The width of the quiresonance region can be classically estimated from the evolution of the phase-space of the perturbed system in the transient process. The angular width W_φ decreases rapidly at low energies and remains nearly constant for large energies, provided that the interaction does not activate high order quiresonances.

We have also analyzed the analogies between the classical and quantum quiresonance behavior. The mayor disagreement between both dynamics occurs at low energies, due to the energy threshold of the quantum diffraction channels. As was described in ultracold atom-diatom collisions [10], no quantitative similarity is found when the inelastic channels are closed. For large translational energies, the quiresonance width coincides, indicating a common mechanism in both dynamics.

Our results for energies up to hundreds of eV indicate an angular quiresonance width $W_\varphi \approx 0.02$ rad, which is above the beam divergence achieved in experimental setups. Although we have assumed a perfect rigid surface, this is a good approximation for large beam energies as the particle wavelength is much smaller than the thermal vibration of the lattice atoms [1]. Therefore, experimental investigations to characterize quiresonance regions in more realistic atom-surface interactions, measuring the diffraction pattern for incident beams closely aligned to the low crystal index direction are in principle possible.

A. R. and J. P. P. acknowledge financial support from Spanish MCT (Grants No. FIS2004-05678, No. FIS2005-02886, and No. FIS2007-64018) and Gobierno de Canarias (Grant No. PI2004/025).

-
- [1] P. Rousseau, H. Khemliche, A. G. Borisov, and P. Rocin, Phys. Rev. Lett. **98**, 016104 (2007).
 - [2] A. Schüller, S. Wethekam, and H. Winter, Phys. Rev. Lett. **98**, 016103 (2007).
 - [3] A. Schüller and H. Winter, Phys. Rev. Lett. **100**, 097602 (2008).
 - [4] D. Gemmell, Rev. Mod. Phys. **46**, 129 (1974).
 - [5] B. Stewart, P. D. Magill, T. P. Scott, J. Derouard, and D. E. Pritchard, Phys. Rev. Lett. **60**, 282 (1988).
 - [6] P. D. Magill, B. Stewart, N. Smith, and D. E. Pritchard, Phys. Rev. Lett. **60**, 1943 (1988).
 - [7] P. D. Magill, T. P. Scott, N. Smith, and D. E. Pritchard, J. Chem. Phys. **90**, 7195 (1989).
 - [8] T. P. Scott, N. Smith, P. D. Magill, D. E. Pritchard, and B. Stewart, J. Phys. Chem. **100**, 7981 (1996).
 - [9] W. J. Hoving and R. Parson, Chem. Phys. Lett. **158**, 222 (1989).
 - [10] R. C. Forrey, N. Balakrishnan, A. Dalgarno, M. R. Haggerty, and E. J. Heller, Phys. Rev. Lett. **82**, 2567 (1999).
 - [11] B. Stewart, P. D. Magill and D. E. Pritchard, J. Phys. Chem. A **104**, 10565 (2000).
 - [12] A. Ruiz and E. J. Heller, Phys. Chem. A **109**, 11578 (2005).
 - [13] A. Ruiz and E. J. Heller, Mol. Phys. **104**, 127 (2006).
 - [14] A. Ruiz, J. P. Palao, and E. J. Heller, J. Phys.: Conf. Ser. **99**, 012018 (2008).
 - [15] A. J. Lichtenberg and M. A. Lieberman, *Regular and Chaotic Dynamics* (Springer-Verlag, New York, 1992).
 - [16] V. Celli, D. Eichenauer, A. Kaufhold, and J. P. Toennies, J. Chem. Phys. **83**, 2504 (1985).